

PESTOFF
... Simply Smarter

RODENT BAIT

RODENT CONTROL SYSTEM

Rodents spoil food, spread disease and cause damage by fouling and gnawing. Although rodents become more visible and annoying in autumn when their numbers are high and they are looking for new homes to escape the winter, sensible rodent control involves year-round preventative action to keep populations in check.

INTRODUCTION

Pestoff introduces Rodent bait, formulated for use indoors or outdoors in approved bait stations, for the control of rats and mice. Brodifacoum (the active ingredient) is a highly effective rodenticide. After two feedings rats will have consumed a lethal dose. Death will usually occur within 4 to 6 days.

MODE OF ACTION

Brodifacoum when regularly fed 2-3 days, cause injury in intestine as well as acting as anti-coagulant. This will cause dehydration, suffocation, anorexia and ischaemia that force rodent to move open space resulting in death ultimately.

PRECAUTIONS

Do not store near / food or feed. Do not lay baits in places accessible to children, pets and domestic animals. Wash hand and exposed skin before meals and after any contact. Burn or bury all rodent bodies and uneaten baits.

If swallowed give a glass or two of water and cause vomiting by putting a finger down the throat. Repeat until vomit fluid is clear in appearance. Vitamin K given orally or intramuscularly is antidotal. In severe causes vitamin K therapy may need to continue for several days or blood transfusions may be necessary.

Manufactured By:

ANIMAL CONTROL PRODUCTS LTD.

408 Heads Road, Wanganui, New Zealand.


Marketed By:

**JAWAD IMPEX
INTERNATIONAL**

210-Rewaz Garden, Lower Mall, Lahore.

Phones: 042-3717 1014, 3717 1145

E-mail: jawadimpexintl@gmail.com



Rodent Blocks

PROCEDURE

- ★ Remove feed from farm vicinity.
 - ★ Wear protecting gloves before handling bait.
 - ★ Keep distance among two bait station 50 ft.
 - ★ First work out rodent load and then decide quantity of bait stations to apply at equal distance.
 - ★ Open lid of box and place two piece of "Pestoff Rodent Block" at bait place.
 - ★ Fix bait stations (pipes / boxes) on path of rodent travel. Bait Station Design (box size 14"x8"x6" made of wood /plastic material having two sides holes or pipe.
 - ★ Regularly check bait station after every 48 hours. Replace bait before eaten by rodent. Continue availability of bait is necessary in bait station.
 - ★ 7-15 days application is sufficient to significantly reduce rodent level in specific area.
 - ★ Apply above action between two flocks (preferably)
- NOTE:** Don't touch bait or bait station with hand because rat has very fast sense of smell against humans.